

## REZZAYO® COMMERCIAL SUSCEPTIBILITY TESTING

<p><b>Hardy Diagnostics</b></p>	<p><b>Kirby Bauer Susceptibility Disk Pack</b>          HardyDisks™ AST          Rezafungin, 5µg, RZF5          Cat. No. Z9471          Pack Size: 1 pack includes          1 cartridge of 50 disks  <a href="https://hardydiagnostics.com/z9471">https://hardydiagnostics.com/z9471</a></p>	<p>HardyDisks™ AST          Rezafungin, 5µg, RZF5          Cat. no. Z9475          Pack Size: 1 pack includes          5 cartridges of 50 disks  <a href="https://hardydiagnostics.com/z9475">https://hardydiagnostics.com/z9475</a></p>
<p><b>ThermoFisher</b></p>	<p><b>Sensititre™ YeastOne™ AST Plate</b>          Sensititre YeastOne IVD Plate,          with Rezafungin          Cat. No. YO4IVD          Pack Size: 10/Pk.  <a href="https://www.thermofisher.com/order/catalog/product/YO4IVD?SID=srch-srp-YO4IVD">https://www.thermofisher.com/order/catalog/product/YO4IVD?SID=srch-srp-YO4IVD</a></p>	<p>Sensititre YeastOne Research Use Only Plate,          w/ Micafungin and Rezafungin          Cat. no. YO11          Pack Size: 10/Pk.  <a href="https://www.thermofisher.com/order/catalog/product/YO11?SID=srch-srp-YO11">https://www.thermofisher.com/order/catalog/product/YO11?SID=srch-srp-YO11</a></p>

## AST METHOD VERIFICATION ISOLATES

<p><b>Laboratory Specialists, Inc.</b></p>	<p><b>Rezafungin (REZZAYO®) – <i>Candida</i> spp.</b>          Cat. No. RZF30          30 strains in set with modal BMD MIC results, based on a minimum of 11 replicates  <a href="https://www.labspec.org/ast-method-verification-isolates-for-new-agents/">https://www.labspec.org/ast-method-verification-isolates-for-new-agents/</a></p>
--	---

## REFERENCE LAB TESTING

<p><b>The Fungal Testing Laboratory</b>          at UT Health San Antonio Long School of Medicine</p>	<p><b>AST Method Providing MIC and Interpretation Susceptibilities</b>  <b>Contact:</b>          Fungus Testing Laboratory Requisition          The University of Texas Health Science Center at San Antonio, Department of Pathology          7703 Floyd Curl Drive, San Antonio, Tx 78229-3900          Tel: (210) 567-4131 Fax: (210) 614-4250          For information on shipping/specimen specific requirements, go to:  <a href="https://lsom.uthscsa.edu/pathology/reference-labs/fungus-testing-laboratory/">https://lsom.uthscsa.edu/pathology/reference-labs/fungus-testing-laboratory/</a></p>
<p><b>ARUP Laboratories</b></p>	<p><b>Fungal Susceptibility Services</b>  <b>Contact:</b>          500 Chipeta Way          Salt Lake City, Utah 84108-1221          Tel: 1-800-522-2787 Fax: 1-800-522-2706          For additional information, please see below or visit:  <a href="https://ltd.aruplab.com/Tests/Pub/2009257">https://ltd.aruplab.com/Tests/Pub/2009257</a></p>
<p><b>Mayo Clinic Laboratories</b></p>	<p><b>AST Method Providing MIC and Interpretation Susceptibilities</b>          Rochester, MN          To Order Rezafungin Test: <b>1.</b> Go to <a href="https://www.mayocliniclabs.com/test-catalog/overview/81602">Mayocliniclabs.com/test-catalog/overview/81602</a>  <b>2.</b> Test ID: MMLYP. <b>3.</b> Click "Order This Test"          Live 24/7 support: 800.533.1710 Fax: 507.284.1759</p>

This information is being provided for reference only and is not an endorsement of any listed product or service. Please contact the listed testing entities for additional information and availability.

**Please see Important Safety Information on reverse side and accompanying Prescribing Information for REZZAYO® (rezafungin for injection).**

# REZZAYO® (rezafungin for injection)

## MEDICAL INFORMATION

For medical inquiries or to report an adverse event, other safety-related information, or product complaints, please contact Medical Information.



1-844-MED-MLNT  
(1-844-633-6568)



medinfo@melinta.com



melintamedicalinformation.com

## INDICATION AND USAGE

REZZAYO® (rezafungin for injection) is an echinocandin antifungal indicated in patients 18 years of age or older who have limited or no alternative options for the treatment of candidemia and invasive candidiasis. Approval of this indication is based on limited clinical safety and efficacy data.

### Limitations of Use

REZZAYO® has not been studied in patients with endocarditis, osteomyelitis, and meningitis due to *Candida*.

## IMPORTANT SAFETY INFORMATION

### Contraindications

REZZAYO® is contraindicated in patients with known hypersensitivity to rezafungin or other echinocandins.

### Warnings and Precautions

- Infusion-related Reactions: REZZAYO® may cause infusion-related reactions, including flushing, sensation of warmth, urticaria, nausea, or chest tightness. If these reactions occur, slow or pause the infusion.
- Photosensitivity: REZZAYO® may cause photosensitivity. Advise patients to use protection from sun exposure and other sources of UV radiation.
- Hepatic Adverse Reactions: Abnormalities in liver tests have been seen in clinical trial patients treated with REZZAYO®. Monitor patients who develop abnormal liver tests and evaluate patients for their risk/benefit of continuing REZZAYO® therapy.

### Adverse Reactions

Most common adverse reactions (incidence  $\geq$  5%) are hypokalemia, pyrexia, diarrhea, anemia, vomiting, nausea, hypomagnesemia, abdominal pain, constipation, and hypophosphatemia.

**Please see accompanying Prescribing Information  
for REZZAYO® (rezafungin for injection).**

All trademarks and registered trademarks are the property of their respective owners.